

ALEXANDROV IMMERSED MINIMAL TORI IN  $S^3$ 

SIMON BRENDLE

ABSTRACT. In this note, we show that our proof of the Lawson Conjecture works for surfaces that are Alexandrov immersed. More precisely, we show that any minimal torus in  $S^3$  which is Alexandrov immersed must be rotationally symmetric. An analogous result holds for surfaces of constant mean curvature.

## 1. INTRODUCTION

In a recent paper [2], we showed the Clifford torus is the only embedded minimal surface in  $S^3$  of genus 1, thereby confirming a conjecture of Lawson. In this note, we classify minimal tori in  $S^3$  that are immersed in the sense of Alexandrov.

**Theorem 1.** *Let  $F : \Sigma \rightarrow S^3$  be an immersed minimal surface in  $S^3$  of genus 1. Moreover, we assume that  $F$  is an Alexandrov immersion; this means that there exists a compact manifold  $N$  and an immersion  $\bar{F} : N \rightarrow S^3$  such that  $\partial N = \Sigma$  and  $\bar{F}|_{\Sigma} = F$ . Then  $\Sigma$  is rotationally symmetric.*

We note that it is possible to classify all rotationally symmetric minimal tori in  $S^3$ ; see [5] for details. This class of surfaces includes the Clifford torus. However, there is a large class of additional examples which are Alexandrov immersed but fail to be embedded.

We will present the proof of Theorem 1 in Section 2. The argument is similar in spirit to the case of embedded surfaces studied in [2], and we will only indicate the necessary modifications.

After the paper [2] was published, Andrews and Li [1] showed that the arguments in [2] can be extended to the setting of constant mean curvature surfaces. As a result, they showed that every embedded constant mean curvature surface in  $S^3$  is rotationally symmetric. Our proof of Theorem 1 also extends to the setting of constant mean curvature surfaces. This yields the following general result:

**Theorem 2.** *Let  $F : \Sigma \rightarrow S^3$  be an immersed constant mean curvature surface in  $S^3$  of genus 1. Suppose that  $F$  extends to an immersion  $\bar{F} : N \rightarrow S^3$  where  $\partial N = \Sigma$  and that  $\partial N$  is mean convex with respect to the pull-back of the standard metric on  $S^3$  under  $\bar{F}$ . Then  $\Sigma$  is rotationally symmetric.*

---

The author was supported in part by the National Science Foundation under grants DMS-0905628 and DMS-1201924.

The proof of Theorem 2 is similar to Theorem 1. The condition that the surface is Alexandrov immersed is quite natural in light of the work of Korevaar, Kusner, and Ratzkin [3] and Kusner, Mazzeo, and Pollack [4], where Alexandrov immersed constant mean curvature surfaces in  $\mathbb{R}^3$  have been studied.

The author is very grateful to Professor Robert Kusner for bringing this problem to his attention, and for many helpful discussions. He would like to thank the Department of Mathematics at Cambridge University, where part of this work was carried out.

## 2. PROOF OF THE MAIN THEOREM

For convenience, we put a Riemannian metric on  $N$  so that  $\bar{F}$  is a local isometry. In particular, there exists a real number  $\delta > 0$  so that  $\bar{F}(x) \neq \bar{F}(y)$  for all points  $x, y \in N$  satisfying  $d_N(x, y) \in (0, \delta)$ .

For each point  $x \in \Sigma$  and any number  $\alpha \geq 1$ , we define

$$D_\alpha(x) = \left\{ p \in S^3 : \frac{\alpha}{\sqrt{2}} |A(x)| (1 - \langle F(x), p \rangle) + \langle \nu(x), p \rangle \leq 0 \right\}.$$

Note that  $D_\alpha(x)$  is a geodesic ball in  $S^3$  whose boundary passes through the point  $F(x)$  and is orthogonal to  $\nu(x)$  at that point.

Let  $I$  denote the set of all points  $(x, \alpha) \in \Sigma \times [1, \infty)$  with the property that there exists a smooth map  $G : D_\alpha(x) \rightarrow N$  such that  $\bar{F} \circ G = \text{id}_{D_\alpha(x)}$  and  $G(F(x)) = x$ .

**Lemma 3.** *Let us fix a pair  $(x, \alpha) \in I$ . Then there is a unique map  $G : D_\alpha(x) \rightarrow N$  such that  $\bar{F} \circ G = \text{id}_{D_\alpha(x)}$  and  $G(F(x)) = x$ .*

**Proof.** Suppose that we can find two distinct maps  $G \neq \tilde{G}$  with these properties. Then  $\bar{F}(G(p)) = \bar{F}(\tilde{G}(p))$  for all points  $p \in D_\alpha(x)$ . This implies  $d_N(G(p), \tilde{G}(p)) \notin (0, \delta)$  for all  $p \in D_\alpha(x)$ . By continuity, we either have  $G(p) = \tilde{G}(p)$  for all  $p \in D_\alpha(x)$  or we have  $G(p) \neq \tilde{G}(p)$  for all  $p \in D_\alpha(x)$ . The second case can be ruled out, as  $G(F(x)) = \tilde{G}(F(x))$ . Thus, we have  $G(p) = \tilde{G}(p)$  for all  $p \in D_\alpha(x)$ . This shows that  $G$  is unique.

**Lemma 4.** *The set  $I$  is closed. Moreover, the map  $G$  depends continuously on the pair  $(x, \alpha)$ .*

**Proof.** Consider a sequence of pairs  $(x^{(m)}, \alpha^{(m)}) \in I$  such that  $\lim_{m \rightarrow \infty} (x^{(m)}, \alpha^{(m)}) = (\hat{x}, \hat{\alpha})$ . For each  $m$ , we can find a smooth map  $G^{(m)} : D_{\alpha^{(m)}}(x^{(m)}) \rightarrow N$  such that  $\bar{F} \circ G^{(m)} = \text{id}_{D_{\alpha^{(m)}}(x^{(m)})}$  and  $G^{(m)}(F(x^{(m)})) = x^{(m)}$ . Since  $\bar{F}$  is a smooth immersion, the maps  $G^{(m)}$  are uniformly bounded in  $C^2$  norm. Hence, after passing to a subsequence, the maps  $G^{(m)}$  converge in  $C^1$  to a map  $G : D_{\hat{\alpha}}(\hat{x}) \rightarrow N$  satisfying  $\bar{F} \circ G = \text{id}_{D_{\hat{\alpha}}(\hat{x})}$  and  $G(F(\hat{x})) = \hat{x}$ . It is easy to see that the map  $G$  is smooth. Thus,  $(\hat{x}, \hat{\alpha}) \in I$ , and the assertion follows.

**Lemma 5.** *We have  $(x, \alpha) \in I$  if  $\alpha$  is sufficiently large.*

**Proof.** By a result of Lawson, we have  $|A(x)| > 0$ . Hence, the radius of the geodesic ball  $D_\alpha(x) \subset S^3$  will be arbitrarily small if  $\alpha$  is sufficiently large. Hence, we can use the implicit function theorem to construct a smooth map  $G : D_\alpha(x) \rightarrow N$  such that  $\bar{F} \circ G = \text{id}_{D_\alpha(x)}$  and  $G(F(x)) = x$ .

After these preparations, we now describe the proof of Theorem 1. Let

$$\kappa = \inf\{\alpha : (x, \alpha) \in I \text{ for all } x \in \Sigma\}.$$

Clearly,  $\kappa \in [1, \infty)$ . For each point  $x \in \Sigma$ , there is a unique map  $G_x : D_\kappa(x) \rightarrow N$  such that  $\bar{F} \circ G_x = \text{id}_{D_\kappa(x)}$  and  $G_x(F(x)) = x$ . For each point  $x \in \Sigma$ , the map  $G_x$  and the map  $\bar{F}|_{G_x(D_\kappa(x))}$  are one-to-one.

We next define a smooth function  $Z : \Sigma \times \Sigma \rightarrow \mathbb{R}$  by

$$Z(x, y) = \frac{\kappa}{\sqrt{2}} |A(x)| (1 - \langle F(x), F(y) \rangle) + \langle \nu(x), F(y) \rangle$$

for  $x, y \in \Sigma$ . In contrast to [2], the function  $Z(x, y)$  might be negative somewhere.

As in [2], we distinguish two cases:

**Case 1:** Suppose first that  $\kappa = 1$ .

**Lemma 6.** *We have  $Z(x, y) \geq 0$  if  $x$  and  $y$  are sufficiently close.*

**Proof.** We argue by contradiction. Suppose that there exist two sequences of points  $x^{(m)}, y^{(m)} \in \Sigma$  such that  $\lim_{m \rightarrow \infty} x^{(m)} = \lim_{m \rightarrow \infty} y^{(m)}$  and  $Z(x^{(m)}, y^{(m)}) < 0$  for all  $m$ . Since  $Z(x^{(m)}, y^{(m)}) < 0$ , the point  $F(y^{(m)})$  lies in the interior of the ball  $D_\kappa(x^{(m)})$ . Therefore, the point  $\tilde{y}^{(m)} := G_{x^{(m)}}(F(y^{(m)}))$  lies in the interior of  $N$ . Since  $y^{(m)}$  lies on the boundary  $\Sigma$ , it follows that

$$\tilde{y}^{(m)} \neq y^{(m)}.$$

On the other hand, we have

$$\bar{F}(\tilde{y}^{(m)}) = F(y^{(m)})$$

and

$$\lim_{m \rightarrow \infty} \tilde{y}^{(m)} = \lim_{m \rightarrow \infty} G_{x^{(m)}}(F(y^{(m)})) = \lim_{m \rightarrow \infty} G_{x^{(m)}}(F(x^{(m)})) = \lim_{m \rightarrow \infty} x^{(m)} = \lim_{m \rightarrow \infty} y^{(m)}.$$

This contradicts the fact that  $\bar{F}$  is an immersion.

As in [2], we perform a Taylor expansion of the function  $Z(x, y)$  when  $x$  and  $y$  are very close. More precisely, let us fix a point  $x \in \Sigma$ , and let  $\{e_1, e_2\}$  is an orthonormal basis of  $T_x \Sigma$  such that  $h(e_1, e_1) > 0$ ,  $h(e_1, e_2) = 0$ , and  $h(e_2, e_2) < 0$ . Let  $\gamma : \mathbb{R} \rightarrow \Sigma$  be a geodesic such that  $\gamma(0) = x$  and  $\gamma'(0) = e_1$ . Since the function  $Z$  is nonnegative when  $x$  and  $y$  are sufficiently close, the function  $f(t) = Z(x, \gamma(t))$  is nonnegative when  $t$  is sufficiently small. Since  $\kappa = 1$ , we have  $f(0) = f'(0) = f''(0) = 0$ . Consequently,  $f'''(0) = 0$ .

This implies that  $(D_{e_1}^\Sigma h)(e_1, e_1) = 0$ . Therefore, the gradient of  $|A|$  at  $x$  is parallel to  $e_2$ . In other words, the function  $|A|$  is constant along one set of curvature lines on  $\Sigma$ . From this, we deduce that  $\Sigma$  is rotationally symmetric.

**Case 2:** Suppose next that  $\kappa > 1$ .

**Lemma 7.** *Fix a point  $x \in \Sigma$ . Then there exists a constant  $\beta > 0$  such that  $d_N(G_x(p), \Sigma) \geq \beta |p - F(x)|^2$  for all points  $p \in \partial D_\kappa(x)$  that are sufficiently close to  $F(x)$ .*

**Proof.** Fix a point  $x \in \Sigma$ . Let us consider the function

$$\varphi_x : \partial D_\kappa(x) \rightarrow \mathbb{R}, p \mapsto d_N(G_x(p), \Sigma).$$

Clearly,  $\varphi_x(F(x)) = 0$ , and the gradient of the function  $\varphi_x$  at the point  $F(x)$  vanishes. Moreover, since  $\kappa > 1$ , the Hessian of the function  $\varphi_x$  at the point  $F(x)$  is positive definite. Hence, we can find a positive constant  $\beta > 0$  such that  $\varphi_x(p) \geq \beta |p - F(x)|^2$  for all points  $p \in \partial D_\kappa(x)$  that are sufficiently close to  $F(x)$ .

**Lemma 8.** *There exists a point  $\hat{x} \in \Sigma$  such that  $\Sigma \cap G_{\hat{x}}(D_\kappa(\hat{x})) \neq \{\hat{x}\}$ .*

**Proof.** Suppose this is false. Then  $\Sigma \cap G_x(D_\kappa(x)) = \{x\}$  for all  $x \in \Sigma$ . This implies that  $d_N(G_x(p), \Sigma) > 0$  for all  $x \in \Sigma$  and all points  $p \in \partial D_\kappa(x) \setminus \{F(x)\}$ . Using the previous lemma, we conclude that there exists a positive constant  $\gamma > 0$  such that  $d_N(G_x(p), \Sigma) \geq \gamma |p - F(x)|^2$  for all  $x \in \Sigma$  and all  $p \in \partial D_\kappa(x)$ . By the implicit function theorem, there exists a small number  $\varepsilon > 0$  such that  $(x, \kappa - \varepsilon) \in I$  for all  $x \in \Sigma$ . This contradicts the definition of  $\kappa$ .

Let  $\hat{x}$  be chosen as in the previous lemma. Moreover, let us pick a point  $\hat{y} \in \Sigma \cap G_{\hat{x}}(D_\kappa(\hat{x}))$  such that  $\hat{x} \neq \hat{y}$ . Since  $\hat{y} \in G_{\hat{x}}(D_\kappa(\hat{x}))$ , we conclude that  $F(\hat{y}) \in D_\kappa(\hat{x})$  and  $G_{\hat{x}}(F(\hat{y})) = \hat{y}$ . Moreover, we claim that  $F(\hat{x}) \neq F(\hat{y})$ ; indeed, if  $F(\hat{x}) = F(\hat{y})$ , then  $\hat{x} = G_{\hat{x}}(F(\hat{x})) = G_{\hat{x}}(F(\hat{y})) = \hat{y}$ , which contradicts our choice of  $\hat{y}$ .

We next consider the function  $Z$  defined above. Since  $F(\hat{y}) \in D_\kappa(\hat{x})$ , we have  $Z(\hat{x}, \hat{y}) \leq 0$ .

**Lemma 9.** *We have  $Z(x, y) \geq 0$  if  $(x, y)$  is sufficiently close to  $(\hat{x}, \hat{y})$ .*

**Proof.** We argue by contradiction. Suppose that there exist sequences of points  $x^{(m)}, y^{(m)} \in \Sigma$  such that  $\lim_{m \rightarrow \infty} x^{(m)} = \hat{x}$ ,  $\lim_{m \rightarrow \infty} y^{(m)} = \hat{y}$ , and  $Z(x^{(m)}, y^{(m)}) < 0$ . Since  $Z(x^{(m)}, y^{(m)}) < 0$ , the point  $F(y^{(m)})$  lies in the interior of the ball  $D_\kappa(x^{(m)})$ . Therefore, the point  $\tilde{y}^{(m)} := G_{x^{(m)}}(F(y^{(m)}))$  lies in the interior of  $N$ . In particular,

$$\tilde{y}^{(m)} \neq y^{(m)}.$$

On the other hand, we have

$$\bar{F}(\tilde{y}^{(m)}) = F(y^{(m)})$$

and

$$\lim_{m \rightarrow \infty} \tilde{y}^{(m)} = \lim_{m \rightarrow \infty} G_{x^{(m)}}(F(y^{(m)})) = G_{\hat{x}}(F(\hat{y})) = \hat{y} = \lim_{m \rightarrow \infty} y^{(m)}.$$

This contradicts the fact that  $\bar{F}$  is an immersion. Thus,  $Z(x, y) \geq 0$  for  $(x, y)$  close to  $(\hat{x}, \hat{y})$ .

Therefore, we can find disjoint open sets  $U, V \subset \Sigma$  such that  $\hat{x} \in U$ ,  $\hat{y} \in V$ , and  $Z(x, y) \geq 0$  for all points  $(x, y) \in U \times V$ . As in [2], we define

$$\Omega = \{x \in U : \text{there exists a point } y \in V \text{ such that } Z(x, y) = 0\}.$$

Since  $Z(\hat{x}, \hat{y}) = 0$ , it follows that  $\hat{x} \in \Omega$ . We can now use the calculation in [2] to conclude that  $Z$  is a supersolution of a degenerate elliptic equation. More precisely, suppose that  $(\bar{x}, \bar{y})$  is an arbitrary point in  $U \times V$ . Then we can find a system of geodesic normal coordinates  $(x_1, x_2)$  around  $\bar{x}$  and a system of geodesic normal coordinates  $(y_1, y_2)$  around  $\bar{y}$  such that

$$\begin{aligned} & \sum_{i=1}^2 \frac{\partial^2 Z}{\partial x_i^2}(\bar{x}, \bar{y}) + 2 \sum_{i=1}^2 \frac{\partial^2 Z}{\partial x_i \partial y_i}(\bar{x}, \bar{y}) + \sum_{i=1}^2 \frac{\partial^2 Z}{\partial y_i^2}(\bar{x}, \bar{y}) \\ & \leq -\frac{\kappa^2 - 1}{\sqrt{2}\kappa} \frac{|A(\bar{x})|}{1 - \langle F(\bar{x}), F(\bar{y}) \rangle} \sum_{i=1}^2 \left\langle \frac{\partial F}{\partial x_i}(\bar{x}), F(\bar{y}) \right\rangle^2 \\ & + \Lambda \left( Z(\bar{x}, \bar{y}) + \sum_{i=1}^2 \left| \frac{\partial Z}{\partial x_i}(\bar{x}, \bar{y}) \right| + \sum_{i=1}^2 \left| \frac{\partial Z}{\partial y_i}(\bar{x}, \bar{y}) \right| \right), \end{aligned}$$

where  $\Lambda$  is a positive constant. Using Bony's version of the strict maximum principle, we conclude that the set  $\Omega$  contains an open neighborhood of  $\hat{x}$ . Moreover, the gradient of  $|A|$  vanishes on the set  $\Omega$ . By analytic continuation,  $|A|$  is a constant function on  $\Sigma$ . This implies that  $F$  is congruent to the Clifford torus. This completes the proof of Theorem 1.

Finally, let us sketch the proof of Theorem 2. Let  $F : \Sigma \rightarrow S^3$  be an immersed constant mean curvature surface in  $S^3$  of genus 1. We assume that  $F$  extends to an immersion  $\bar{F} : N \rightarrow S^3$  where  $\partial N = \Sigma$  and that  $\partial N$  is mean convex with respect to the pull-back of the standard metric on  $S^3$  under  $\bar{F}$ . Given a point  $x \in \Sigma$  and a real number  $\alpha \geq 1$ , one defines

$$D_\alpha(x) = \left\{ p \in S^3 : \left( \frac{H}{2} + \frac{\alpha}{\sqrt{2}} |\mathring{A}(x)| \right) (1 - \langle F(x), p \rangle) + \langle \nu(x), p \rangle \leq 0 \right\}$$

(cf. [1]). Here,  $H$  is the mean curvature (i.e. the sum of the principal curvatures) and  $\mathring{A}$  is the trace-free part of the second fundamental form. As above, let  $I$  denote the set of all points  $(x, \alpha) \in \Sigma \times [1, \infty)$  with the property that there exists a smooth map  $G : D_\alpha(x) \rightarrow N$  such that  $\bar{F} \circ G = \text{id}_{D_\alpha(x)}$  and  $G(F(x)) = x$ . Finally, let

$$\kappa = \inf \{ \alpha : (x, \alpha) \in I \text{ for all } x \in \Sigma \}.$$

Combining the arguments above with the calculations in [1] and [2], one concludes that  $F$  is rotationally symmetric.

#### REFERENCES

- [1] B. Andrews and H. Li, *Embedded constant mean curvature tori in the three-sphere*, preprint (2012)
- [2] S. Brendle, *Embedded minimal tori in  $S^3$  and the Lawson conjecture*, to appear in Acta Mathematica
- [3] N. Korevaar, R. Kusner, and J. Ratzkin, *On the nondegeneracy of constant mean curvature surfaces*, Geom. Funct. Anal. 16, 891–923 (2006)
- [4] R. Kusner, R. Mazzeo, and D. Pollack, *The moduli space of complete embedded constant mean curvature surfaces*, Geom. Funct. Anal. 6, 120–137 (1996)
- [5] O. Perdomo, *Rotational surfaces in  $S^3$  with constant mean curvature*, preprint (2012)

DEPARTMENT OF MATHEMATICS, STANFORD UNIVERSITY, STANFORD, CA 94305